

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS21 4677

Anti-GST-tag (mouse, monoclonal)

## **Product information**

**Immunogen** GST (glutathione S-transferase) recombinant protein

Host Mouse

Clonality Monoclonal

Subclass/isotype IgG1a

**Purity** Affinity purified in PBS pH 7.4. Contains 0.02 % sodium azide. Contains 50 % glycerol.

Format Liquid

**Quantity** 50 g

Store at -20 C for up to 3 years, Make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the

tube.

## **Application information**

Recommended dilution 1: 500 - 1: 5000 (WB)

Expected | apparent

Depends upon MW of a fusion protein

**Additional information** This antibody is recognizing native and denatured forms of GST

Proteins were isolated from E. coli Rosetta cells transformed with pGEX6P-1 vector. Lane 1 contains protein fraction from bacterial cells before induction with IPTG (-). Lane 2 contains protein fraction from bacterial cells after 2 h induction with IPTG (+). 15uL of sample were loaded and separated on an 15 % SDS-PAGE and transferred for 70 min at 55V using a tank transfer system to nitrocellulose membrane. Blots were blocked with 1x PBS + 0.1 %Tween 20 (T) + 5% milk for 1 h at room temperature (RT) with agitation. Blots were incubated with the primary antibody -GST (AS21 4677) at indicated dilutions overnight at 4 C with agitation in 1x PBS-T + 5% milk. The primary antibody solutions were decanted, and the blots were washed 4 times (6-8 minutes each) in 1x PBS-T at RT with agitation. Blots were incubated with secondary antibody Goat anti-Mouse (Product no: AS09 627-trial) diluted to 1: 20,000 in 1x PBS-T + 5% milk for 2 h at RT with agitation. The blots were washed as above and developed with ECL or chemiluminescent detection reagents AgriseraECL SuperBright (AS16 ECL-S-10) for 4 min and imaged using ChemiDoc (BioRad) for 5 seconds. PonceauS served as loading control. Courtesy of Kelly Mason and Antje Heese; Dept. Biochemistry, IPG; University of Missouri, Columbia (MO), USA