


Product no **AS23 4914****Anti-PFK1-7 | Phosphofructokinase 1-7****Product information**

<b>Immunogen</b>	KLH-conjugated peptide derived from <i>Arabidopsis thaliana</i> PFK1-7 protein sequences, UniProt: <a href="#">Q9M0F9</a> , <a href="#">Q9FIK0</a> , <a href="#">Q94AA4</a> , <a href="#">Q9FKG3</a> , <a href="#">Q8VYN6</a> , <a href="#">Q9M076</a> , <a href="#">Q9C5J7</a> TAIR: At4g29220, At5g47810, At4g26270, At5g61580, At2g22480, At4g32840, At5g56630
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Antigen affinity purified serum, in PBS pH 7.4
<b>Format</b>	Lyophilized
<b>Quantity</b>	50 µg
<b>Reconstitution</b>	For reconstitution add 50 µl, of sterile or deionized water.
<b>Storage</b>	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tubes.

**Application information****Recommended dilution** | 1 : 1000 - 1 : 2000 (WB)**Expected | apparent MW** | 51.9 kDa**Confirmed reactivity** | *Arabidopsis thaliana*

**Predicted reactivity** | *Brassica napus* (PFK1, PFK3, PFK5, PFK6, PFK7),  
*Nicotiana tabacum* (PFK3, PFK4, PFK5, PFK6)  
*Pisum sativum* (PFK3, PFK4, PFK5, PFK7, PFK6)  
*Solanum lycopersicum* ( PFK3, PFK4, PFK5, PFK6)  
*Solanum tuberosum* ( PFK3, PFK4, PFK5, PFK6)

*Arachis hypogaea*, *Brachypodium distachyon*, *Brassica napus*, *Cannabis sativa*, *Hordeum vulgare*, *Malus domestica*, *Manihot esculenta*, *Medicago truncatula*, *Nicotiana tabacu*, *Oryza sativa*, *Saccharum sp.*, *Theobroma cacao*, *Triticum sp.*, *Sorghum bicolor*, *Zea mays*, *Vitis vinifera*

Species of your interest not listed? [Contact us](#)**Not reactive in** | No confirmed exceptions from predicted reactivity are currently known**Selected references** | To be added when available, antibody available in January 2025.

Samples:

20 µg of *Arabidopsis thaliana* whole leaf extract20 µg of *Arabidopsis thaliana* mutant described above the picture MW markers are marked on the left side of each membrane

20 µg/well of total protein extracted freshly from leaf extracts of *Arabidopsis thaliana* wildtype and mutants. Exact buffer components were: 50 mM Hepes-KOH (pH 6.8), 5 mM Mg-acetate, 15 % Glycerin, 1 mM EDTA, 1 mM EGTA, 5 mM β-Mercaptoethanol, 0.1 mM Pefabloc Proteinase-inhibitor and denatured with 1 x Laemmli-buffer (62.5 mM Tris-HCl (pH 6.8), 2 % SDS, 10 % Glycerin, 5 % β-Mercaptoethanol, 0.001 % Bromophenolblue) at 98 °C / 2 mins. Samples were separated on 10 % SDS-PAGE and blotted for 1 h PVDF (pore size of 0.45 µm), using: wet transfer in the cold. Blot was blocked with 5 % nonfat milk 4°C / ON with agitation. Blot was incubated in the primary antibody at a dilution of 1: 2 000 for 1 h/RT with agitation in TBS-T. The antibody solution was decanted, and the blot was rinsed briefly, then washed once 3 times for 10 min

This product is **for research use only** (not for diagnostic or therapeutic use)

contact: [support@agrisera.com](mailto:support@agrisera.com)

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | [www.agrisera.com](http://www.agrisera.com)

in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1: 25 000 in for 1 h/RT with agitation. The blot was washed as above and developed with a following chemiluminescent detection reagent. Exposure time was 3 minutes.

Courtesy of phd student Alina Johanna Hieber, University of Bayreuth, Bayreuth, Germany