

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS21 4528 Anti-PEC1 | Plastid Envelope Channel 1

Product information

Immunogen	The soluble domain 466 (M244 until stop codon, 64 kDa) was cloned into pET16b and transformed into BLR 21 for 467 expression in <i>Escherichia coli</i> UniProt: <u>Q8VZM7-1</u> , TAIR: <u>At5g02940</u>
Host	Rabbit
Clonality	Polycional
Purity	Serum
Format	Lyophilized
Quantity	50 µl
Reconstitution	For reconstitution add 50 μ l, of sterile water
Storage	Store lyophilized/reconstituted at -20°C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please remember to spin the tubes briefly prior to opening them to avoid any losses that might occur from material adhering to the cap or sides of the tube.
nal information	This antibody is recognizing PEC1, but not PEC2 or DMI1

Application information

Additio

Recommended dilution	1 : 1000 (WB)
Expected apparent MW	100 75 kDa
Confirmed reactivity	Arabidopsis thaliana
Predicted reactivity	Camelina sativa, Capsella rubella, Nicotiana benthamiana, Noccaea caerulescens, Pisum sativum, Raphanus sativus,
Selected references	Species of your interest not listed? <u>Contact us</u> <u>Völkner</u> et al (2021) Two plastid POLLUX ion channel-like proteins are required for stress-triggered stromal Ca2+release, Plant Physiology, Volume 187, Issue 4, December 2021, Pages 2110–2125,
	https://doi.org/10.1093/plphys/kjab424



Samples:

- 1-2,5 ug of Arabidopsis thaliana mutant pec1-1pec2-1
- 2-2,5 ug of Arabidopsis thaliana mutant pec1-2pec2-2
- 3-2,5 ug of Arabidopsis thaliana whole leaf extract

Arabidopsis thaliana tissue was frozen with liquid nitrogen and ground to a fine powder with mortar and pestle. Protein was extracted by mixing with extraction buffer (200 mM tris pH 8.0, 4% (w/v) sodium dodecyl sulfate) to 0.5 g fresh weight/mL and heating at 80 °C for 8 min. 5 µl protein extract were separated on 10 % SDS-PAGE and blotted at 75v for 45min to nitrocellulose (pore size of 20 um), using wet transfer. Blot was blocked with 5% milk for 1h/RT with agitation. Blot was incubated in the primary antibody at a dilution of 1: 1000 ON/4 °C in 5% Milk-TBS-T with agitation. The antibody solution was decanted and the blot was rinsed briefly twice, then washed once for 15 min and 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in matching secondary antibody (anti-rabbit IgG horse radish peroxidase conjugated) diluted to 1:10 000 in for 1h/RT with agitation. The blot was washed as above and developed for 1 min with chemiluminescent detection reagent following manufacture's recommendations. Exposure time was 120 seconds.

Courtesy of Carsten Voelkner, Kunz Lab